

NON-PERTURBATIVE EFFECTIVE LAGRANGIANS FOR SUPER-MATRIX MODELS

Ram Brustein^{*)}

Theoretical Physics Division, CERN
CH - 1211 Geneva 23

and

Dept. of Physics, Univ. of Pennsylvania
Philadelphia, PA 19104, U.S.A.

Michael Faux^{*)} and Burt A. Ovrut^{),***)}**

Theoretical Physics Division, CERN
CH - 1211 Geneva 23

ABSTRACT

We discuss $d = 1, \mathcal{N} = 2$ supersymmetric matrix models and exhibit the associated $d = 2$ collective field theory in the limit of dense eigenvalues. From this field theory we construct, by the addition of several new fields, a $d = 2$ supersymmetric effective field theory, which reduces to the collective field theory when the new fields are replaced with their vacuum expectation values. This effective theory is Poincare invariant and contains perturbative and non-perturbative information about the associated superstrings.

^{*)} Work supported in part by DOE under Contract No. DOE-AC02-76-ERO-3071.

^{**)} On sabbatical leave from the Department of Physics, University of Pennsylvania, Philadelphia, PA 19104, U.S.A.

^{***)} Invited talk, International Europhysics Conference on High Energy Physics, Marseille, France, July 22-28, 1993.

The question of how supersymmetry is broken remains one of the most important issues in superstring theory. The advent of $d = 1$ supersymmetric matrix models[1], which contain nonperturbative information about a class of two-dimensional superstrings, offers the hope that one can identify, and study quantitatively, the exact string mechanism for supersymmetry breaking. To do this, it is essential to construct the complete two-dimensional supersymmetric effective Lagrangian associated with this class of theories. Once that is achieved, one can look for non-perturbative phenomena, such as instantons, that will induce supersymmetry violating operators (spontaneous or explicit) in this effective Lagrangian. This entire program has already been carried out[2], with positive results, in the case of $d = 1$ bosonic matrix models where the nonperturbative effective Lagrangian was constructed and its fundamental symmetry, a non-compact shift symmetry, $\zeta \rightarrow \zeta + c$ in one of its fields ζ , was shown to be broken by instantons in a single eigenvalue of the matrix model. In this talk we will discuss the extension of these results to the supersymmetric case[3]. To be precise, we will construct the two-dimensional supersymmetric effective Lagrangian for the superstrings associated with the $d = 1$ supersymmetric matrix model. We have also discovered instantons in the eigenvalues of these matrix models, which have all the right characteristics to break supersymmetry. However, the constraints placed on the length of this manuscript mandate that we discuss these instantons and supersymmetry breaking elsewhere.

The simplest supersymmetric matrix model which admits non-trivial interactions involves a $d = 1, \mathcal{N} = 2$ supersymmetry. The fundamental variable of the theory is a time-dependent $N \times N$, Hermitian matrix superfield

$$\Phi_{ij} = M_{ij}(t) + i\theta_1 \Psi_{1ij}(t) + i\theta_2 \Psi_{2ij} + i\theta_1 \theta_2 F_{ij}(t), \quad (1)$$

where θ_1 and θ_2 are the two anticommuting parameters. The Lagrangian is given by

$$L = \int d\theta_1 d\theta_2 \left\{ \frac{1}{2} \text{Tr} D_1 \Phi D_2 \Phi + iW(\Phi) \right\}, \quad (2)$$

where the superpotential is $W(\Phi) = \sum_n b_n \text{Tr} \Phi^n$. Eliminating the auxiliary fields F_{ij} , quantizing the theory and restricting the Hilbert space to the $U(N)$ invariant states, one arrives[4, 3] at an effective component field Lagrangian of the form

$$\begin{aligned} L = & \sum_i \left\{ \frac{1}{2} \dot{\lambda}_i^2 - \frac{1}{2} \left(\frac{\partial W_{eff}(\lambda)}{\partial \lambda_i} \right)^2 \right\} \\ & - \frac{i}{2} \sum_i (\psi_{1i} \dot{\psi}_{1i} + \psi_{2i} \dot{\psi}_{2i}) - i \sum_{ij} \psi_{1i} \frac{\partial^2 W_{eff}(\lambda)}{\partial \lambda_i \partial \lambda_j} \psi_{2j}. \end{aligned} \quad (3)$$

where $W_{eff} = W + w$ and $w(\lambda) = -\sum_i \sum_{j \neq i} \ln |\lambda_i - \lambda_j|$. The λ_i are the N eigenvalues of M_{ij} and ψ_i are associated fermions. Lagrangian (3) is $d = 1, \mathcal{N} = 2$ supersymmetric. This theory can be converted from one dimensional quantum mechanics to a two-dimensional field theory by introducing a real parameter, x , restricted to the interval $-L/2 < x < L/2$ and by defining the collective fields

$$\begin{aligned} \varphi(x, t) &= \sum_i \Theta(x - \lambda_i(t)) \\ \psi(x, t) &= -\sum_i \delta(x - \lambda_i(t)) \chi_i(t). \end{aligned} \quad (4)$$

The limit of physical interest is when $N \rightarrow \infty, L \rightarrow \infty$. See[5] for previous work on this subject. If $N/L \rightarrow \infty$, then the eigenvalues become dense in L and the collective fields (4) become unconstrained space-time fields. In this case, rewriting (3) in terms of the collective fields and appropriately taking the $N \rightarrow \infty, L \rightarrow \infty$ limit, one finds that the Lagrangian becomes

$$L = \int dx \left\{ \frac{\dot{\varphi}^2}{2\varphi'} - \frac{\pi^2}{6}\varphi'^3 + \frac{1}{2}\omega^2 x^2 \varphi' - \frac{i}{2\varphi'}(\psi_1 \dot{\psi}_1 + \psi_2 \dot{\psi}_2) - \frac{i\pi}{2}\psi_1 \psi'_1 + \frac{i\pi}{2}\psi_2 \psi'_2 + \frac{i}{2}\frac{\dot{\varphi}}{\varphi'^2}(\psi_1 \psi'_1 + \psi_2 \psi'_2) \right\}, \quad (5)$$

where $\psi = \frac{1}{\sqrt{2}}(\psi_1 + i\psi_2)$ and ω^2 is an arbitrary positive constant. The equations of motion associated with (5) have a static solution with $\langle \psi_1 \rangle = \langle \psi_2 \rangle = 0$ and $\langle \varphi' \rangle = \frac{1}{\pi} \sqrt{\omega^2 x^2 - \frac{1}{g}}$, where g is an arbitrary positive integration constant. In this case, $\langle \varphi \rangle$ is only defined in the region $|x| \geq 1/\omega\sqrt{g}$. Therefore, in this region only, we can expand φ as $\varphi = \langle \varphi \rangle + \frac{1}{\sqrt{\pi}}\zeta$. Furthermore, if we define a new spatial coordinate τ through the expression $\tau' = \frac{1}{\pi} \langle \varphi' \rangle^{-1}$, the Lagrangian (5) can be reexpressed as

$$L = \int d\tau \left\{ \frac{1}{2}(\dot{\zeta}^2 - \zeta'^2) - \frac{1}{2} \frac{g(\tau)\dot{\zeta}^2 \zeta'}{1 + g(\tau)\zeta'} - \frac{1}{6}g(\tau)\zeta'^3 + \frac{1}{3g(\tau)^2} - \frac{i}{\sqrt{2}}(\psi_+ \dot{\psi}_+ - \psi_+ \psi'_+) - \frac{i}{\sqrt{2}}(\psi_- \dot{\psi}_- + \psi_- \psi'_-) + \frac{i}{\sqrt{2}} \frac{g(\tau)\zeta'}{1 + g(\tau)\zeta'}(\psi_+ \dot{\psi}_+ + \psi_- \dot{\psi}_-) + \frac{i}{\sqrt{2}} \frac{g(\tau)\dot{\zeta}}{(1 + g(\tau)\zeta')^2}(\psi_+ \psi'_+ + \psi_- \psi'_-) \right\}, \quad (6)$$

where $\psi_{+,-} = \frac{2^{1/4}}{\sqrt{\pi}}\psi_{1,2}$, $g(\tau)$ is a space dependant coupling parameter given by

$$g(\tau) = \frac{4\sqrt{\pi}e^{-2\omega(\tau-\tau_0)}}{\omega(1 - \frac{1}{\omega g}e^{-2\omega(\tau-\tau_0)})^2}. \quad (7)$$

and τ_0 is an arbitrary constant. Note that the Lagrangian (6) suffers from the fact that the coupling parameter $g(\tau)$ is explicitly a function of τ and, hence, L is not Poincare invariant. To overcome this difficulty, we make the assumption that $g(\tau)$ arises as the vacuum expectation value (VEV) of a function of a new scalar field, which we denote by α . We infer the existence of an effective Poincare invariant theory involving α , as well as ζ, ψ_+ and ψ_- , which reproduces (6) when α is replaced by its τ -dependant VEV. Additionally, we postulate that this effective field theory possesses a $d = 2$ supersymmetry. It follows that, in addition to α , we must introduce its fermionic superpartners χ_+ and χ_- which, of course, have vanishing VEV'S. There are an infinite number of $d = 2, (p, q)$ supersymmetries. It is not hard to demonstrate that the

appropriate supersymmetry for our effective field theory is (1,1) supersymmetry. The original and newly introduced fields discussed above appear as components of the (1,1) superfields

$$\begin{aligned}\Phi_1 &= \zeta + i\theta^+\psi_+ + i\theta^-\psi_- + i\theta^+\theta^-Z \\ \Phi_2 &= \alpha + i\theta^+\chi_+ + i\theta^-\chi_- + i\theta^+\theta^-A.\end{aligned}\quad (8)$$

The construction of the correct supersymmetric effective field theory proceeds as follows. Consider the free part of the collective field Lagrangian. It is given by

$$\mathcal{L}_{01} = \frac{1}{2}(\dot{\zeta}^2 - \zeta'^2) - i\psi_+\partial_-\psi_+ - i\psi_-\partial_+\psi_-.\quad (9)$$

Now consider the manifestly supersymmetric Lagrangian

$$\begin{aligned}\mathcal{L}_{01}^{(eff)} &= \int d\theta^+d\theta^-D_+\Phi_1D_-\Phi_1 \\ &= \frac{1}{2}(\dot{\zeta}^2 - \zeta'^2) - i\psi_+\partial_-\psi_+ - i\psi_-\partial_+\psi_- + Z^2\end{aligned}\quad (10)$$

Field Z is auxiliary and can be eliminated using its equation of motion, $\langle Z \rangle = 0$. It is clear that (10) reproduces (9). We also need to introduce a kinetic energy for Φ_2 . To do this, we must find a superfield Lagrangian involving Φ_2 only, whose component field equations admit $\langle \alpha \rangle = \exp(-\omega(\tau - \tau_0))$, $\langle \chi_{\pm} \rangle = \langle A \rangle = 0$ as a solution, and which reduces to the $1/3g(\tau)^2$ term in (6) when these VEV's are inserted. The appropriate Lagrangian is

$$\begin{aligned}\mathcal{L}_{02}^{(eff)} &= \int d\theta^+d\theta^- \left\{ F_1(\Phi_2)D_+\Phi_2D_-\Phi_2 \right. \\ &\quad \left. - \frac{1}{\omega^2}F_2(\Phi_2)\partial_-D_+\Phi_2\partial_+D_-\Phi_2 \right\}\end{aligned}\quad (11)$$

where F_1 and F_2 are polynomial functions of Φ_2 whose exact form is irrelevant for this discussion. We proceed to study the interaction terms in (6). The terms in (6) linear in the coupling $g(\tau)$ are

$$\begin{aligned}\mathcal{L}_1 &= g(\tau) \left\{ -\frac{1}{6}(\zeta'^3 + 3\dot{\zeta}^2\zeta') \right. \\ &\quad \left. + \frac{i}{\sqrt{2}}\zeta'(\psi_+\dot{\psi}_+ + \psi_-\dot{\psi}_-) + \frac{i}{\sqrt{2}}\dot{\zeta}(\psi_+\psi'_+ + \psi_-\psi'_-) \right\}.\end{aligned}\quad (12)$$

We would like to find a superfield Lagrangian involving Φ_1 and Φ_2 which, when added to (10) and (11), continues to admit $\langle \alpha \rangle = \exp(-\omega(\tau - \tau_0))$, $\langle \chi_{\pm} \rangle = \langle A \rangle = 0$ and $\langle Z \rangle = 0$ as a solution to the equations of motion and which, when these are substituted, reduces to (12). We have shown that the appropriate Lagrangian is

$$\begin{aligned}\mathcal{L}_1^{(eff)} &= \int d\theta^+d\theta^- \left\{ \right. \\ &\quad \frac{f(\Phi_2)}{\omega^3\Phi_2^3}\partial_{(+}\Phi_1\partial_{-)}\Phi_2\partial_{[+}\Phi_1\partial_{-]}\Phi_2D_{(+}\Phi_1D_{-)}\Phi_2 \\ &\quad - \frac{f(\Phi_2)}{\omega^5\Phi_2^5}(\partial_{[+}\Phi_1\partial_{-]}\Phi_2)^2\partial_{(+}\Phi_1\partial_{-)}\Phi_2D_+\Phi_2D_-\Phi_2 \\ &\quad \left. + \frac{1}{3}\frac{f(\Phi_2)}{\omega^5\Phi_2^5}(\partial_{(+}\Phi_1\partial_{-)}\Phi_2)^3D_+\Phi_2D_-\Phi_2 \right\},\end{aligned}\quad (13)$$

where

$$f(\Phi_2) = \frac{4\sqrt{\pi}}{\omega} \frac{\Phi_2^2}{(1 - \frac{1}{g\omega}\Phi_2^2)^2}, \quad (14)$$

$A_{(+B_-)} = A_+B_- + A_-B_+$, and $A_{[+B_-]} = A_+B_- - A_-B_+$. It is possible to extend this expression to all orders in $g(\tau)$,

$$\begin{aligned} \mathcal{L}^{(eff)} = \int d\theta_+ d\theta_- \bigg\{ & D_+ \Phi_1 D_- \Phi_1 \\ & + F_1(\Phi_2) D_+ \Phi_2 D_- \Phi_2 - \frac{1}{\omega^2} F_2(\Phi_2) \partial_- D_+ \Phi_2 \partial_+ D_- \Phi_2 \\ & + \frac{f(\Phi_2)}{\omega^3 \Phi_2^3} \frac{\partial_{(+}\Phi_1 \partial_{-)}\Phi_2 \partial_{[+}\Phi_1 \partial_{-]}\Phi_2}{1 + \frac{f(\Phi_2)}{\omega\Phi_2} \partial_{(+}\Phi_1 \partial_{-)}\Phi_2} D_{(+}\Phi_1 D_{-)}\Phi_2 \\ & - \frac{f(\Phi_2)}{\omega^5 \Phi_2^5} \frac{(\partial_{[+}\Phi_1 \partial_{-]}\Phi_2)^2 \partial_{(+}\Phi_1 \partial_{-)}\Phi_2}{1 + \frac{f(\Phi_2)}{\omega\Phi_2} \partial_{(+}\Phi_1 \partial_{-)}\Phi_2} D_+ \Phi_2 D_- \Phi_2 \\ & + \frac{1}{3} \frac{f(\Phi_2)}{\omega^5 \Phi_2^5} (\partial_{[+}\Phi_1 \partial_{-]}\Phi_2)^3 D_+ \Phi_2 D_- \Phi_2 \bigg\}. \end{aligned} \quad (15)$$

The component field equations of motion have $\langle \alpha \rangle = \exp(-\omega(\tau - \tau_0))$, $\langle \chi_{\pm} \rangle = \langle A \rangle = 0$ and $\langle Z \rangle = 0$ as a solution. When these are substituted back into the component field expansion of (15), one obtains exactly the collective field Lagrangian, (6). Therefore, the Lagrangian (15) is the two-dimensional supersymmetric effective Lagrangian for the superstrings associated with the $d = 1$ supersymmetric matrix model.

References

- [1] E. Marinari and G. Parisi, *Phys. Lett.* **B247** (1990) 537.
- [2] R. Brustein and B. Ovrut, *Phys. Lett.* **B309** (1993) 45 and preprint, UPR-523T (1992).
- [3] R. Brustein, M. Faux and B. Ovrut, CERN-TH. 7017 (1993).
- [4] A. Dabholkar, *Nucl. Phys.* **B368** (1992) 293.
- [5] A. Jevicki and J. Rodrigues, *Phys. Lett.* **B268** (1991) 53; J. Rodrigues and J. van Tonder, preprint, CNLS-92-02 (1992); J. Cohn and H. Dykstra, *Mod. Phys. Lett.* **A7** (1992) 1163; J. Feinberg, preprint, TECHNION-PH-92-35 (1992).